

**Patients having surgery in the morning:**

- **Do not** have Sodium-glucose co-transporter-2 (SGLT2) inhibitors (see list in the table below) for **72 hours** before surgery.
- **Do not** have other diabetic tablets the evening before or the morning of procedure. Bring all diabetic medications with you.
- **Have normal evening dose and half morning dose** of subcutaneous insulin. Exanatide/Liraglutide/Dulaglutide - brand name Byetta/Bydureon or Victoza or Trulicity.

**Patients having surgery in the afternoon:**

- **Do not** have Sodium-glucose co-transporter-2 (SGLT2) inhibitors (see list in the table below) for **72 hours** before surgery.
- **Do not** have other diabetic tablets the evening before or the morning of procedure, even if having breakfast. Bring all diabetic medications with you.
- **Have normal evening dose and half morning dose** of subcutaneous insulin. Exanatide/Liraglutide/Dulaglutide - brand name Byetta/Bydureon or Victoza or Trulicity.

<b>Class</b>	<b>Generic (Brand) name</b>
Biguanides	• Metformin • Metformin XR
Sulfonylureas	• Glibenclamide • Gliclazide • Gliclazide MR • Glimepiride • Glipizide
Dipeptidyl peptidase-4 (DPP-4) inhibitors	• Alogliptin • Linagliptin • Saxagliptin • Sitagliptin • Vildagliptin
Thiazolidinediones (TZD)	• Pioglitazone • Rosiglitazone
Glucagon-like peptide-1 (GLP-1) receptor agonists	• Dulaglutide • Exenatide • Exenatide ER • Liraglutide • Lixisenatide
Sodium-glucose co-transporter-2 (SGLT2) inhibitors	• Empagliflozin + Linagliptin (Glyxambi) • Empagliflozin + Metformin (Jardiamet) • Dapagliflozin + Saxagliptin (Qtern) • Dapagliflozin + Metformin (Xigduo) • Ertugliflozin + Sitagliptin (Steglujan) • Ertugliflozin + Metformin (Segluromet)

Note: This **list is not exhaustive** so please check if you are on a medication for diabetes.

